Blue Cross and Blue Shield of Rhode Island Preventive Services for Commercial Members Jan 2021 Note: See separate policy for Preventive Services for Blue CHIP for Medicare Members Coverage for newly recommended preventive health services is made no later than one year after the release date of the recommendation. Although many USPSTF recommendations are gender specific, BCBSRI does not limit coverage for gender specific services that are **denotes an updated recommendation **denotes an updated recommendation **denotes an updated recommendation Unhealthy Alcohol Use in Adolescents and Adults USPSTF Grade: B (Nov 2018) The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use. Unhealthy Drug Use: Screening USPSTF Grade: B (June 2020)** The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.) Health Service Code(s) Frequency			
Dowerage for newly recommended preventive health services is made no later than one year after the release date of the recommendation.			
Inhough many USPSTF recommendations are gender specific, BCBSRI does not limit coverage for gender specific services that are denotes an updated recommendation **denotes a new recommendation **denotes an ewer commendation **denotes an updated recommendation **denotes an ewer commendation **denotes an ewer comme			
denotes an updated recommendation **denotes a new recommendation nhealthy Alcohol Use in Adolescents and Adults USPSTF Grade: B (Nov 2018) the USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant omen and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy cohol use. nhealthy Drug Use: Screening USPSTF Grade: B (June 2020)** the USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be uplemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers asking questions about unhealthy drug use, not testing biological specimens.)			
nhealthy Alcohol Use in Adolescents and Adults USPSTF Grade: B (Nov 2018) be USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant omen and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy cohol use. hhealthy Drug Use: Screening USPSTF Grade: B (June 2020)** be USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be plemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers asking questions about unhealthy drug use, not testing biological specimens.)			
he USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant omen and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy cohol use. Inhealthy Drug Use: Screening USPSTF Grade: B (June 2020)** The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be applemented when services for accurate diagnosis; effective treatment, and appropriate care can be offered or referred. (Screening refers a saking questions about unhealthy drug use, not testing biological specimens.)			
The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant owner and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy lookol use. Inhealthy Drug Use: Screening USPSTF Grade: B (June 2020)**			
omen and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy cohol use. Inhealthy Drug Use: Screening USPSTF Grade: B (June 2020)** The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be mplemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers a saking questions about unhealthy drug use, not testing biological specimens.)			
inhealthy Drug Use: Screening USPSTF Grade: B (June 2020)** The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be mplemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers a saking questions about unhealthy drug use, not testing biological specimens.)			
The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be mplemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers o asking questions about unhealthy drug use, not testing biological specimens.)			
The USPTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be mplemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers o asking questions about unhealthy drug use, not testing biological specimens.)			_
mplemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers o asking questions about unhealthy drug use, not testing biological specimens.)			
o asking questions about unhealthy drug use, not testing biological specimens.)			
99408, 99409 No Specific Code 1 per year			
99385-99387 99395-99397 No Specific Code 2 per year any combination of codes			
9/1029/			
	l.		
sychosocial/Behavioral Assessments for ages newborn to age 21 Bright Futures/American Academy of Pediatrics			
Health Service Code(s) Diagnosis Code(s) Frequency			
99381-99384 No Specific Code 2 per year any combination of codes			
77,371-77,374			
epression Screening			$ \top$
~Adolescents USPSTF Grade: B (February 2016) he USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be		_	-+
ne USF31F recomments screening for major depressive disorder (MDD) in adoescents aged 12 to 18 years. Screening should be inplemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up.			
right Futures/American Academy of Pediatrics			-+
Health Service Code(s) Diagnosis Code(s) Frequency			
99384-99385 No Specific Code 2 per year any combination of codes			
99394-99395			
96127 No Specific Code None (no annual limit) -Adults USPSTF Grade: B (January 2016)			
he USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women.			
creening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate			
llow up. Health Service Code Diagnosis Code Frequency			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes 99395-99397 No Specific Code None (no annual limit)			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes 99395-99397 No Specific Code None (no annual limit) Idernal Depression Screening terening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". AP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months.			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes 99395-99397 No Specific Code None (no annual limit) faternal Depression Screening creening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". AP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. right Futures/American Academy of Pediatrics			
Health Service Code Diagnosis Code Frequency 9383-99387 99395-99397 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) Internal Depression Screening recenting should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". AP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. right Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency			
Health Service Code Diagnosis Code Frequency 93835-99387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) aternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Per recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency None None (no annual limit)			
Health Service Code Diagnosis Code Frequency 9385-9387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) aternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None (no annual limit) parately reimbursed up to 12 months of			
Health Service Code Diagnosis Code Frequency 93835-93887 No Specific Code 2 per year any combination of codes 93935-93937 No Specific Code None (no annual limit) atternal Depression Screening treening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". AP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Flutures/American Academy of Pediatrics Health Service Code 96161 None None None (no annual limit) parately reimbursed up to 12 months of e; Over 12 months covered and not			
Health Service Code Diagnosis Code Frequency 9383-99387 99395-99397 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) Internal Depression Screening recenting should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". AP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. right Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes 90385-99387 No Specific Code None (no annual limit) aternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Percommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. giph Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None None (no annual limit) parately reimbursed up to 12 months of g: Over 12 months covered and not			
Health Service Code Diagnosis Code Frequency 93835-9387 No Specific Code 2 per year any combination of codes 93935-93937 No Specific Code None (no annual limit) aternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Percommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None (no annual limit) parately reimbursed up to 12 months of e; Over 12 months covered and not			
Health Service Code Diagnosis Code Prequency 93835-93837 No Specific Code 2 per year any combination of codes 93935-93937 No Specific Code None (no annual limit) atternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". PAP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Prequency 96161 None None (no annual limit) parately reimbursed up to 12 months of eye (over 12 months covered and not parately reimbursed over 12 months covered and not parately reimbursed			
Health Service Code Diagnosis Code Prequency 93835-93887 No Specific Code 2 per year any combination of codes 93935-93937 No Specific Code None (no annual limit) atternal Depression Screening treening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Per recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code 96161 None Prequency 96161 None None (no annual limit) parately reimbursed up to 12 months of e; Over 12 months covered and not parately reimbursed parately reimbursed the USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal the USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal przession to counseling interventions.			
Health Service Code Piagnosis Code Piagnosis Code Prequency 9385-99387 No Specific Code Pol 2 per year any combination of codes 96127 No Specific Code None (no annual limit) atternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code None None None None (no annual limit) Practice of the property of t			
Health Service Code 93855-93887 No Specific Code 2 per year any combination of codes 95955-93877 No Specific Code None (no annual limit) aternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends spediatrician academy of Pediatrics Health Service Code Diagnosis Code Diagnosis Code Frequency None None (no annual limit) None (no annual limit) Every specific Code None (no annual limit) None (no annual limit) Parately reimbursed up to 12 months of export of the president of the pre			
Health Service Code Diagnosis Code Prequency 93835-99387 No Specific Code 2 per year any combination of codes 93935-99397 No Specific Code None (no annual limit) atternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". APP recommends spediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None (no annual limit) parately reimbursed up to 12 months of eye (over 12 months covered and not parately reimbursed of the commend			
Health Service Code Diagnosis Code Prequency 93385-93887 No Specific Code 2 per year any combination of codes 95395-9397 No Specific Code None (no annual limit) aternal Depression Screening aternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". PAP recommends spediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None (no annual limit) Parately reimbursed up to 12 months of export of the present			
Health Service Code Diagnosis Code Prequency 99385-99387 No Specific Code 2 per year any combination of codes 99385-99387 No Specific Code None (no annual limit) aternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". AP recommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None None (no annual limit) parately reimbursed up to 12 months of et. Over 12 months covered and not be pression to counseling interventions. Health Service Code Diagnosis Code(s) Diagnosis Code(s) Frequency Population of the USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Health Service Code(s) Diagnosis Code(s) Frequency Population (covered and not separately reimbursed) None (no annual limit) None (no annual limit)			
Health Service Code Diagnosis Code Prequency 99385-99387 No Specific Code 2 per year any combination of codes 90385-99387 No Specific Code None (no annual limit) aternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Prequency 96161 None None None (no annual limit) parately reimbursed up to 12 months of e; Over 12 months covered and not parately reimbursed Intellection of the present of the present of perinatal persons who are at increased risk of perinatal pression to counseling interventions. Health Service Code(s) Diagnosis Code(s) Frequency 10 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			
Health Service Code 93835-9387 No Specific Code 2 per year any combination of codes 93935-93937 No Specific Code None (no annual limit) atternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code None None None (no annual limit) Poarately reimbursed up to 12 months of experiments covered and not be contactly reimbursed or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Health Service Code(s) Diagnosis Code(s) Prequency None (no annual limit) Pression to counseling interventions. Health Service Code(s) Podd(covered and not separately reimbursed); 99401 (covered and not separately reimbursed); 99402-99404 No Specific Code No Specific Code None (no annual limit) No Specific Code None (no annual limit)			
Health Service Code Diagnosis Code Prequency 99385-99387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) Maternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Percommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None None (no annual limit) Pression Into Pediatric Practice". When the Management of Perinatal Depression Into Pediatric Practice			
Health Service Code Diagnosis Code 2 per year any combination of codes 93835-93837 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) atternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Percommends pediatrican screening during newborn visits at I month, 2 months, 4 months and 6 months. Ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None None (no annual limit) Pression to counseling interventions of experiments covered and not be acately reimbursed pression to counseling interventions. Health Service Code(s) Diagnosis Code(s) Frequency 19401 (covered and not separately reimbursed): No Specific Code None (no annual limit)			
Health Service Code Diagnosis Code Prequency 99385-99387 No Specific Code 2 per year any combination of codes 99395-99397 No Specific Code None (no annual limit) Maternal Depression Screening Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Ight Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Frequency 96161 None None None (no annual limit) Marately reimbursed up to 12 months of C; Over 12 months covered and not warately reimbursed WEPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Health Service Code Diagnosis Code(s) Frequency 99401 (covered and not separately reimbursed): No Specific Code None (no annual limit) No Specific Code None (no annual limit) 99411-99412 (covered but not separately reimbursed): 99384-99386 No Specific Code 2 per year any combination of codes			
Health Service Code Diagnosis Code Frequency 9385-99387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) Internal Depression Screening Recenting should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Precommends for the service Code Diagnosis Code Frequency Precommends for no annual limit) Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Prequency Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Prequency Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Prequency Precommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Prequency Precommends that clinicians provide or refer pregnant and postpa			
Health Service Code Diagnosis Code Frequency 9385-99387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) teternal Depression Screening recening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at 1 month, 2 months, 4 months and 6 months. By Frequency Prequency 96161 None None (no annual limit) parately reimbursed up to 12 months of (Over 12 months covered and not arrately reimbursed pression to counseling interventions. Health Service Code(s) Diagnosis Code(s) Frequency 99401 (covered and not separately reimbursed) 99401 (covered and not separately reimbursed) 99402-99404 No Specific Code None (no annual limit) 90411-99412 (covered but not separately reimbursed) 90384-99386 No Specific Code 2 per year any combination of codes 90791, 90792 Z13.32, and pregnancy diagnosis codes O00.00-O9A.53 None (no annual limit) velopmental and Autism Screening Pregnancy Precommends Prequency P			
Heath Service Code Diagnosis Code Frequency 9935-99397 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) Internal Depression Screening Internal Depression Into Pediatric Practice*. Precommends pediatrican screening during newborn visits at 1 month, 2 months, 4 months and 6 months. Internal Depression Into Pediatric Practice*. Prequency 96161 None Prequency 96161 None None (no annual limit) Internal Service Code Diagnosis Code Prequency Internal Depression Into Pediatric Practice*. Internal Depression Into Pediatrice*. Internal Depression Into Pediatrice Practice*. Internal Depression Into Pediatrice*. Internal Depression Into Pediatrice			
Health Service Code Diagnosis Code Frequency 9335-99387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) None (no annual			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes 99385-99397 No Specific Code None (no annual limit) Internal Depression Screening Verification Ver			
Health Service Code Diagnosis Code Frequency 9335-99387 No Specific Code 2 per year any combination of codes 96127 No Specific Code None (no annual limit) None (no			
Health Service Code 99385-99387 No Specific Code 2 per year any combination of codes 99385-99397 No Specific Code None (no annual limit) Internal Depression Screening Internal Depression Into Pediatric Practice". Internal Depression Screening Internal Depression Into Pediatric Practice". Internal Depression Internal Inter			
Health Service Code Diagnosis Code Frequency 99385-99387 No Specific Code 2 per year any combination of codes 99385-99397 No Specific Code None (no annual limit) atternal Depression Sercening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatricias recenting during newborn visits at I month, 2 months, 4 months and 6 months. ght Futures/American Academy of Pediatrics Health Service Code Diagnosis Code Prequency 90161 None None (no annual limit) ice USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal pression to counseling interventions. Health Service Code(s) Diagnosis Code(s) Frequency 99401 (covered and not separately reimbursed) No Specific Code None (no annual limit) 99401-99401 No Specific Code None (no annual limit) 99401-99402-99404 No Specific Code None (no annual limit) 99411-99412 (covered but not separately reimbursed) 99384-99396 No Specific Code 2 per year any combination of codes 99394-99396 No Specific Code 2 per year any combination of codes 99394-99396 No Specific Code 2 per year any combination of codes 99394-99396 No Specific Code None (no annual limit) 99401-99401 None (no annual limit) 99402-99403 None (no annual limit) 99403-9936 No Specific Code 2 per year any combination of codes 99394-99396 None (no annual limit) 99404-99409 None (no annual limit)			
Health Service Code Diagnosis Code Prequency 93935-99397 No Specific Code 2 per year any combination of codes 99395-99397 No Specific Code None (no annual limit)			
Health Service Code 97938-599387 No Specific Code 1 None (no annual limit) sternal Depression Screening reening should occur per "Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice". Precommends pediatrician screening during newborn visits at I month, 2 months, 4 months and 6 months. ### Futures/American Academy of Pediatrics Health Service Code Diagnosis Code None None (no annual limit) #### Supplied or refer pregnant and postpartum persons who are at increased risk of perinatal persons to counseling interventions. ###################################			
Health Service Code 993859387 No Specific Code 12 per year any combination of codes 9939599397 No Specific Code None (no annual limit) 16127 No Specific Code None (no annual limit) 1628 1628 1629 1629 1629 1629 1629 1629 1629 1629			
Health Service Code 97935-97937 No Specific Code 1 2 per year any combination of codes 97935-97937 No Specific Code None (no annual limit) 1 1			
Health Service Code 979385-97987 No Specific Code 1 Prequency 979395-97997 No Specific Code 1 None (no annual limit) 1 None (no annual limit) 1 None (no annual limit) 1 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 1 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 1 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 1 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 2 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 3 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 4 Precommends pediatrician screening during newborn visits at 1 month, 2 months and 6 months. 5 Prequency 6 1611 None None (no annual limit) 1 Precommends during the covered and not arrachy reimbursed 2 Prequency 2 Prequency 3 Prequency 4 Prequency 4 Prequency 5 Prequency 5 Prequency 6 Prequency 6 Prequency 6 Prequency 6 Prequency 6 None (no annual limit) 7 Prequency 7 Prequency 7 Prequency 8 No Specific Code None (no annual limit) 7 Prequency 7 Prequency 8 No Specific Code None (no annual limit) 7 Prequency 8 No Specific Code None (no annual limit) 8 Prequency 9 Prequency 9 Prequency 9 Prequency 9 Prequency 9 Prequency 1 Prequency 2 Prequency 2 Prequency 3 Prequency 4 Prequency 4 Prequency 4 Prequency 5 Prequency 5 Prequency 6 Prequency 7 Prequency			
Health Service Code 97935-97937 No Specific Code 1 2 per year any combination of codes 97935-97937 No Specific Code None (no annual limit) 1 1			